

Homework IV
41000 - Business Statistics
Spring 2013

Due dates: 05/20(41000-81) - 05/21(41000-82)

Turn in your homework at the beginning of the class. Your homework solution should be well organized. Include graphs that you are asked to view in the problems. You are not limited or restricted to use the statistical packages used/presented by the instructor. You may discuss the problems with your colleagues but in the end you MUST write up your homework individually.

Problem I: Let us check the claim that the daily closing returns of Apple Inc. (AAPL) and Microsoft Corporation (MSFT) in 2009 are just as likely to be positive or negative. Assume that, day to day, positive returns are all independent Bernoulli(p), where p is the probability of positive return. Of the 217 business days in 2009, AAPL was positive for 124 days, while MSFT was positive for 120 days.

- a) Draw the Binomial(217,0.5) distribution. This is the distribution of the number of days out of 217 with positive returns should the probability of positive return for a given day be 0.5. Use the excel function BINOMDIST.
- b) Locate the observed data on the graph, i.e. 124 days in AAPL and 120 days in MSFT;
- c) Obtain 95% confidence intervals for p_A ;
- d) Obtain 95% confidence intervals for p_M ;
- e) Test the hypothesis $H_0: p_A = 0.5$ for AAPL at the 5% level;
- f) Test the hypothesis $H_0: p_M = 0.5$ for MSFT at the 5% level;
- g) Compute the P-value for e);
- h) Compute the P-value for f).

Problem II: Let x_1, \dots, x_{1000} be sample A and y_1, \dots, y_{1000} be sample B (see Figure 2 below).

- a) Two statisticians, Shapiro and Wilk, devised in 1965 a statistical procedure to test normality of a data set. I implemented the Shapiro-Wilk's normality test for both samples, i.e. I tested H_0 : "data is normal". The P-values are equal to 0.0000000000426 and 0.2917, respectively. What can we say about the assumption of normality for the data sets?
- b) Two other statisticians, Durbin and Watson, devised in 1950 a statistical procedure to test serial correlation of a data set. If there is serial correlation then today's data is similar to yesterday's data and so forth. I implemented the Durbin-Watson's serial correlation test for both samples, i.e. we tested H_0 : "data is not serially correlated". The P-values are to 0.08906 and 0.4622, respectively. What can we say about the assumption of no serial correlation for the data sets?




Figure 2: Sample A and sample B.